









Identification of Carbon Stars from LAMOST DR7

Linlin Li¹ , Kecheng Zhang¹, Wenyuan Cui¹ , Jianrong Shi² , Wei Ji^{3,4} , Zhenyan Huo¹, Yawei Gao¹, Shuai Zhang¹ , and Mingxu Sun¹ 

¹ Department of Physics, Hebei Normal University, Shijiazhuang 050024, People's Republic of China; wenyuancui@126.com

² Key Lab of Optical Astronomy, National Astronomical Observatories, Chinese Academy of Sciences, A20 Datun Road, Chaoyang, Beijing 100101, People's Republic of China

³ Key Laboratory of Space Astronomy and Technology, National Astronomical Observatories, CAS, Beijing 100101, People's Republic of China

⁴ School of Astronomy and Space Science, University of Chinese Academy of Sciences, Beijing 100049, People's Republic of China

Received 2023 August 4; revised 2023 November 22; accepted 2023 December 4; published 2024 February 20

Speaker: Bao Yuxi

Date : 2024/04/26

Outline

- 1 Background
- 2 Data
- 3 Identification and Spectral Classification
- 4 Discussion and Conclusion

1 Background

- Carbon stars are excellent kinematic tracers of galaxies and play important roles in understanding the evolution of the Galaxy.
- Carbon stars are rare and peculiar objects first discovered by Secchi (1869), and characterized by strong carbon molecular bands, such as CH, CN, and C₂, in their optical spectra.
- Carbon stars usually have similar temperature and luminosity with late G, K, and M stars, while showing higher carbon abundance than oxygen ($C/O > 1$; Keenan 1993).
- According to their different spectral features, carbon stars can be classified into different subclasses (Keenan 1993; Barnbaum et al. 1996), including C–H, C–R, C–N, C–J, and Ba stars.
- They can also be divided into giant carbon and dC (dwarf Carbon) stars depending on their brightness.

1 Background

- Almost all Ba and C–H stars are binaries (McClure et al. 1980; McClure 1983; Jorissen et al. 2016; Escorza & De Rosa 2023; Goswami & Goswami 2023).
- Aikman & McClure (1984) carried out a series of radial velocity monitoring for C–N stars and found signs of binary companions for C–N stars.
- However, McClure (1997) did not find any evidence of binary for 22 C–R stars after 16 yr of radial velocity monitoring.

2 Data

- LAMOST can simultaneously obtain the spectra of 4000 targets in a single exposure, and its resolution is ~ 1800 for the low-resolution mode.
- The DR7 data set contains 14,487,406 spectra, including 10,599,979 low-resolution spectra, and 1,008,710 non-time-domain and 2,878,717 time-domain medium-resolution spectra.
-
- In the LAMOST DR7 Low-Resolution Spectroscopic (LRS) General Catalog, there are 4402 spectra of 3565 stars labeled as carbon for the “subclass” parameter (stellar spectral type).
- However, the contamination rate is not presented, and no classification of the carbon stars is given.
- In this work, we systematically search for different types of carbon stars from more than 10 million low-resolution spectra.

2.1 Line Indices

- They use the multiple [line index](#) to select carbon stars, as the spectra of carbon stars are characterized by strong absorption bands of carbon-containing molecules.
- The line index is defined by the following equation according to the equivalent width (Worthey et al. 1994; Liu et al. 2015):

$$EW = \int \left[1 - \frac{f_{\text{line}}(\lambda)}{f_{\text{cont}}(\lambda)} \right] d\lambda, \tag{1}$$

where $f_{\text{cont}}(\lambda)$ and $f_{\text{line}}(\lambda)$ are the fluxes of the [continuum](#) and the [spectral line](#).

The line index under this definition is in Å.

We measure [five line indices](#) including C₂(λ 4737 Å, λ 5165 Å, and λ 5635 Å) and CN (λ 7065 Å, and λ 7820 Å). The definition of line indices is listed in Table 1.

Table 1
Line Indices Definition in This Work

Name (1)	Index Bandpass (Å) (2)	Pseudocontinua (Å) (3)
C ₂ (4737 Å)	4620–4742	4580–4620 4742–4812
C ₂ (5165 Å)	4980–5170	4930–4980 5170–5235
C ₂ (5635 Å)	5350–5640	5300–5350 5640–5700
CN(7065 Å)	7065–7190	7025–7065 7190–7230
CN(7820 Å)	7820–8000	7790–7820 8000–8040

3 Identification and Spectral Classification

➤ 3.1 The Identification Steps of Carbon Stars

- The [high S/N spectra](#) are helpful to us to identify and classify carbon stars more accurately.

Different types of stars are mixed, and it is very difficult to identify the carbon stars in the highly contaminated region.





Figure 1. Distribution of the 9,060,920 spectra.

➤ 3.1 The Identification Steps of Carbon Stars

- They further remove pollution based on the infrared color– color diagram.







Figure 2. Distribution of the 706,597 candidates and the known carbon stars in the J- H vs. J- Ks color–color diagram. The solid dots indicate the density of the candidates.

➤ 3.1 The Identification Steps of Carbon Stars



The lower left of the dashed line is the Warm Group, the upper right is the Cool Group.



The red dashed line: Si et al. (2015)

Figure 3. Distribution of the 609,702 candidates and the known carbon stars in the $J-H$ vs. $H-K_s$ color-color diagram.

➤ The Warm Group





Figure 4. Distribution of the 586,076 warm candidates and the known carbon stars in the $EW_{C_2(\lambda 5635 \text{ \AA})}$ vs. $EW_{C_2(\lambda 4737 \text{ \AA})}$ plane.




Figure 5. Distribution of the 127,943 warm candidates and the known carbon stars in the $EW_{C_2(\lambda 5635 \text{ \AA})}$ vs. $EW_{CN(\lambda 7065 \text{ \AA})}$ plane.

➤ The Cool Group





Figure 6. Distribution of 23,629 cool candidates and known carbon stars in the $EW_{C_2}(\lambda 5635 \text{ Å})$ vs. $EW_{CN}(\lambda 7820 \text{ Å})$ plane.

➤ 3.2 Spectral Classification

- For the spectral classification of carbon stars in this work, we adopt the latest carbon star classification system published by Barnbaum et al. (1996).

1: C–H Stars

- Strong **G band** at $\lambda 4300$ Å.
- Secondary **P-branch head** near $\lambda 4342$ Å as a distinctive feature.
- Weak **Fe I line** at $\lambda 4383$ Å.
- Strong **Ba II lines** at $\lambda 4554$ Å and $\lambda 6496$ Å.

2: C–R Stars:

- Strong **CN absorption band** at $\lambda 4215$ Å.
- Prominent **Ca I line** at $\lambda 4226$ Å.
- Very weak **Ba II lines** at $\lambda 4554$ Å and $\lambda 6496$ Å.

3: C–N Stars:

- Almost **no flux** below $\lambda 4400$ Å, or even $\lambda 5000$ Å, a key identifier.
- Isotopic bands are generally weak.
- Enhanced lines of **s-process elements**, especially **Ba**, more than in C–R stars.

4: Ba Stars:

- Significant enhancement** of s-process elements.
- Strong **Ba II line** at $\lambda 4554$ Å and **Sr II line** at $\lambda 4077$ Å.




Figure 7. Four typical LAMOST spectra of carbon stars classified.

➤ 3.3 Identification Results

Table 2
Carbon Stars in This Work

Subtype	Number of Carbon Stars	Number of Carbon Stars Spectra
C-H	925	1217
C-R	284	341
C-N	608	799
Ba	1292	1657
UNKNOWN	437	528
Total Number	3546	4542

They manually classify 4542 spectra of 3546 carbon stars identified in LAMOST DR7.




Figure 9. Steps of the identification of carbon stars.

4.1 Distribution of Carbon Stars

- JHKs colors are a good indicator of effective temperature (Wang & Jiang 2014); therefore, they are often used to distinguish subtype carbon stars with different temperatures.




Figure 10. Distribution of our carbon stars in the J – H vs. H– K_s diagram.

- Previous works have shown that many carbon stars are actually carbon dwarf stars (Green 2013; Green et al. 2019; Roulston et al. 2022).





Figure 11. Distribution of our carbon stars in the H-R diagram.

4.2 Spatial Distribution



The majority of C–N (95%) and a C–R (72%) and Ba stars (84%) are concentrated at **low Galactic latitudes**.

A majority of the C–H stars and dC stars (70.9%) lie in **high Galactic latitudes**.

dC stars are from the older thick-disk and halo populations (Farihi et al. 2018; Roulston et al. 2022).

Figure 13. Spatial distribution of the 3546 carbon stars reported in this paper in the Galactic coordinates.

Summary

- From over 10 million low-resolution spectra in LAMOST DR7, we identified 3546 carbon stars (4542 spectra), including 925: C–H, 1392: Ba, 608: C–N, and 284: C–R stars, with 437: unclassified carbon stars marked as "UNKNOWN".
- By mapping carbon star candidates in the H-R diagram, we identified 258 dwarf carbon star candidates from the 3546: carbon stars, comprising 33: Ba, 104: C–H, 21: C–R, and 100: "UNKNOWN" stars.
- We discussed the spatial distribution of each subtype of carbon star in the Milky Way, finding that most carbon stars are distributed in the anti-galactic direction, related to LAMOST's sky survey strategy. The spatial distribution confirms that C–H stars are primarily found in halo populations, while C–N, C–R, and Ba stars are mostly distributed at low Galactic latitudes. Among the 437 "UNKNOWN" stars, about 40% are located in regions with $|b| \geq 30^\circ$.

Thanks